arXiv:1207.1050v1 [astro-ph.CO] 4 Jul 2012

Is a co-rotating Dark Disk a threat to Dark Matter Directional Detection ?

J. Billard,* Q. Riffard, F. Mayet, and D. Santos
Laboratoire de Physique Subatomique et de Cosmologie,
Université Joseph Fourier Grenoble 1, CNRS/IN2P3,
Institut Polytechnique de Grenoble, Grenoble, France
(Dated: July 5, 2012)

Recent N-Body simulations are in favor of the presence of a co-rotating Dark Disk that might contribute significantly (10%-50%) to the local Dark Matter density. Such substructure could have dramatic effect on directional detection. Indeed, in the case of a null lag velocity, one expects an isotropic WIMP velocity distribution arising from the Dark Disk contribution, which might weaken the strong angular signature expected in directional detection. For a wide range of Dark Disk parameters, we evaluate in this Letter the effect of such dark component on the discovery potential of upcoming directional detectors. As a conclusion of our study, using only the angular distribution of nuclear recoils, we show that Dark Disk models as suggested by recent N-Body simulations will not affect significantly the Dark Matter reach of directional detection, even in extreme configurations.

PACS numbers: 95.35.+d, 14.80.-j

I. INTRODUCTION

Within the standard Dark Matter halo paradigm, the local Dark Matter distribution is assumed to be smoothly spatially distributed and to be well-described by a Maxwellian velocity distribution. However, the hierarchical structure formation model indicates that the Galactic Dark Matter halo results from sucessive small halo accretions, thus directly linking its structure to its merging history. The presence of substructures in the Milky Way halo is inferred from recent results of N-body simulation [1–6]. Such substructures may be classified as follows : Dark Matter tidal streams (spatially localized), debris flows (spatially homogenized but with velocity substructures) and a Dark Disk. The latter has received much interest since late sub-halo merging is expected to lead to the formation of a co-rotating Dark Disk [6–9] that may affect the expected WIMP signal both in direct and directional detection. While the influence of the dark disk on Dark Matter signals has been exhaustively investigated for direct [6, 10–12] and indirect [13] detection, it is still unclear how it may affect directional detection. Following a previous work from A. M. Green [12], we aim at evaluating the influence of the presence of a co-rotating Dark Disk on the discovery potential of a forthcoming directional detector. In particular, we are interested in determining the values of the Dark Disk parameters for it to significantly affect the Dark Matter reach of directional detection. In order to be model independent from the background energy modelling, the study has been done by considering only the angular distribution of nuclear recoils $dR/d\Omega_r$.

Since the pionner paper of D. N. Spergel [14], the contribution of directional detection to the field of Dark Matter has been adressed through a wealth of studies [4, 15–36]. Depending on the unknown WIMP-nucleon cross section, directional detection may be used to : exclude Dark Matter [15, 16], reject the isotropy hypothesis [17–23], discover galactic Dark Matter with a high significance [24–26] or constrain WIMP and halo properties [27–29]. For neutralino Dark Matter, a large fraction MSSM configurations with a neutralino lighter than 200 GeV/c^2 would lead to a significance greater than 3σ (90% CL) in a 30 kg.year CF_4 directional detector [30].

In the following, we focus on the effect of a co-rotating Dark Disk on the potential of forthcoming directional detectors to discover Dark Matter [26]. The paper is organized as follows. Section II presents the current knowledge on the Dark Disk. In particular, we define its parametrization used throughout. The directional framework is recalled in sec. III, with emphasize on the directional statistic methods used to exploit forthcoming data. Then, for a wide range of Dark Disk parameters, we evaluate in IV the effect of such substructure on the discovery potential of upcoming directional detectors.

II. A DARK DISK IN THE MILKY WAY

Recent results from N-body simulation of Milky Way type galaxies have shown that merging satellite galaxies may get dragged into the plane of their host galaxy [6–9]. This leads to a Dark Matter overdensity roughly matching the baryonic disk of the host galaxy and usually in co-rotation with the latter [7, 8]. This Dark Matter component is usually referred to as Dark Disk (DD). Hitherto, there is no observational evidence in favor of a Dark Disk in the Milky Way. In fact, a Dark Disk density greater than 50% of the halo one ($\rho_{DD}/\rho_H > 0.5$) is excluded by an estimation of the dynamical surface mass density in the solar neighbourhood obtained from the kinematics of a sample of 1200 red giant stars [37].

The Dark Disk is generally considered as a cold substructure for which the velocity distribution is decribed by an

---

*Electronic address: billard@lpsc.in2p3.fr

isotropic Maxwellian distribution [7]. In such context, the astrophysical parameters relevant to the description of the Dark Disk are the density ρ_{DD}, its co-rotational velocity V_{DD} and its velocity dispersion given by σ_{DD}. The range of interest of these parameters must be inferred from the results of N-body simulations and compared to astrophysical constraints. For instance, F. S. Ling *et al.* [6] have extracted a Milky way type galaxy from the results of the RAMSES simulation [1]. The velocity distribution of Dark Matter particles within a $7 < R < 9$ kpc and $|Z| < 1$ kpc is shown to be well fitted by a double gaussian along the ϕ direction. The first one, corresponding to the Dark Matter halo component, is described by a null average speed and a velocity dispersion $\sigma_{halo} \simeq 180$ km/s. The second component, the Dark Disk, is described by a co-rotation velocity $V_{DD} \simeq 150$ km/s and a velocity dispersion $\sigma_{DD} \simeq 85$ km/s. Note that this description is in good agreement with [7–9]. However, as there is no clear observational constraints on these parameters, we allow for a wide range in order to investigate the effect of a Dark Disk component on directional detection. Unless otherwise stated, we consider hereafter the following Dark Disk parameter ranges:

$$\begin{array}{rcl} 0 < & \rho_{DD}/\rho h & < 1 \\ 0\ \mathrm{km.s^{-1}} < & V_{DD} & < 220\ \mathrm{km.s^{-1}} \\ 7\ \mathrm{km.s^{-1}} < & \sigma_{DD} & < 155\ \mathrm{km.s^{-1}} \end{array} \quad (1)$$

III. DIRECTIONAL DETECTION FRAMEWORK

There is a worldwide effort toward the development of a large TPC (Time Projection Chamber) devoted to directional detection [38–42]. In the following, we exemplify our Dark Disk study considering a MIMAC-like detector corresponding to a low exposure (30 kg.year) CF_4 TPC allowing three dimensional track measurements [41].

The two dimensional directional recoil rate $d^2R/dE_r d\Omega_r$ is given by [35] :

$$\frac{\mathrm{d}^2R}{\mathrm{d}E_r\mathrm{d}\Omega_r} = \frac{\rho_0\sigma_0}{4\pi m_\chi m_r^2} F^2(E_r)\hat{f}(v_{\min},\hat{q}), \quad (2)$$

with m_χ the WIMP mass, m_r the WIMP-nucleus reduced mass, ρ_0 the local Dark Matter density, σ_0 the WIMP-nucleus elastic scattering cross section, $F(E_r)$ the form factor (using the axial expression from [43]), $v_{\min}$ the minimal WIMP velocity required to produce a nuclear recoil of energy E_r and $\hat{q}$ the direction of the recoil momentum. Finally, $\hat{f}(v_{\min},\hat{q})$ is the three-dimensional Radon transform of the WIMP velocity distribution $f(\vec{v})$. As the Radon transform is a linear application, one can simply add the host halo and the Dark Disk contribution to the directional event rate. The angular distribution $dR/d\Omega_r$ is thus obtained by integrating the double-differential spectrum over the energy range chosen to be $E_r = [5, 50]$ keV.

There are several approaches to exploit the forthcoming directional data. Either the data analysis may aim at rejecting the isotropy hypothesis [17–23] or at discovering galactic Dark Matter with a high significance [26] via a profile likelihood ratio test. In the following, we investigate the effect of a Dark Disk contribution on the expected significance of a discovery of Dark Matter with directional detection. To do so, we use a twofold approach, following [20, 26].

First, we use a profile likelihood ratio test statistic, as presented in [26] and briefly recalled hereafter for the reader's convenience. In order to remain independent from the background energy spectrum modelling, only the directional information is considered in the following, *i.e.* the angular distribution of the recoiling events $dR/d\Omega_r$. Noting σ_p the WIMP-proton cross section and R_b the background rate, the likelihood function is given by,

$$\begin{aligned} \mathscr{L}(\sigma_p, R_b) = {} & \frac{(\mu_s+\mu_b)^N}{N!}e^{-(\mu_s+\mu_b)} \\ & \times \prod_{n=1}^{N}\left[\frac{\mu_s}{\mu_s+\mu_b}S(\vec{R}_n) + \frac{\mu_b}{\mu_s+\mu_b}B(\vec{R}_n)\right] \end{aligned} \quad (3)$$

where $\mu_b = R_b \times \xi$ and μ_s corresponds to the number of expected background and WIMP events respectively, where ξ corresponds to the exposure. N is the total number of observed events, $\vec{R}_n$ refers to the direction of each event while the functions S and B are the directional event rate $dR/d\Omega_r$ of the WIMP and background events respectively. Following recent studies on the angular distribution of muon induced neutrons [44], the background angular distribution B is assumed to be isotropic in the galactic rest frame. Note that, contrary to a previous work [26] the astrophysical uncertainties are not taken into account in the estimation of the significance to allow fair comparison between the two statistical approaches. Only the background rate is taken as a nuisance parameter.

In a frequentist approach, the significance of a new process is commonly estimated by using the profile likelihood ratio test [45]. It corresponds to a hypothesis test of the null hypothesis H_0 (background only) against the alternative H_1 which includes both background and signal. As discussed in [45] the test statistic in the case of a discovery is defined as follows:

$$\mathrm{q}_0 = \begin{cases} -2\ln\lambda(0) & \hat{\sigma_p} > 0 \\ 0 & \hat{\sigma_p} < 0 \end{cases} \quad (4)$$

with,

$$\lambda(0) = \frac{\mathscr{L}(\sigma_p = 0, \hat{\hat{R}}_b)}{\mathscr{L}(\hat{\sigma_p}, \hat{R}_b)} \quad (5)$$



FIG. 1: Mean significance E(Z) as a function of V_{DD}, the rotation velocity of the Dark Disk at Solar radius. Left : the result is presented for various values relative density ρ_{DD}/ρ_H and a fixed velocity dispersion $\sigma_{DD} = 85$ km/s. Right : the result is presented for various values of the velocity dispersion σ^{DD} and a fixed value of the relative density $\rho_{DD}/\rho_H = 0.5$. These studies has been done for 50 GeV/c^2 WIMP mass and a fixed value of 100 WIMP events (from the halo and the Dark Disk).

Hence, a large value of q_0 implies a large discrepancy between the two hypothesis which is in favor of a discovery (H_1). As $f(q_0 \mid H_0)$ follows a χ^2_1 distribution, the discovery significance Z is simply defined as $Z = \sqrt{q^{\rm obs}}$, in units of σ [45].

The second approach, first introduced by B. Morgan *et al.* [20], is based on a generic test of isotropy following the mean recoil deviation $\langle \cos\theta \rangle$ such as:

$$\langle \cos\theta \rangle = \frac{1}{N}\sum_{i=1}^{N} \cos\theta_i \qquad (6)$$

where θ_i is the i^{th} angle between the recoil and the Cygnus direction, and N is the number of measured recoils. Note that this test is, by definition, coordinate system dependent, such as the previous one as the main recoil direction (ℓ, b) (see [25]) is not considered here as a fitting parameter.

Eventually, one can evaluate the significance of an observed anisotropy by computing the distributions of $\langle \cos\theta \rangle$ for both H_0 corresponding to the background (isotropic) and H_1 the alternative. It is worth noticing that the use of the variable $\langle \cos\theta \rangle$ is particularly interesting in the case of directional detection of Dark Matter as the expected signal should exhibit a dipole feature hence maximizing the deviation between H_0 and H_1.

IV. INFLUENCE OF A CO-ROTATING DARK DISK

In order to investigate the effect of a Dark Disk component on the expected significance of a directional dark matter detection, we allow for a wide range on the Dark Disk parameters, see eq. 1, and we evaluate, for each configuration, the expected significance for a 30 kg.year MIMAC-like detector. We highlight the fact that for a co-rotating Dark Disk to contribute to the data, the energy threshold must be low and/or the WIMP mass large. For concreteness, we present a case study for a 50 GeV/c^2 WIMP mass and a total of 100 WIMP events. Figure 1 (left) presents the mean significance E(Z) as a function of V_{DD}, the rotation velocity of the Dark Disk at Solar radius. The black dashed line corresponds to the no Dark Disk case. The result is then presented for various values of the relative density ρ_{DD}/ρ_H. The general feature is that the mean significance is decreasing when increasing the co-rotating velocity of the Dark Disk at Solar radius as it results in a loss of directionality. This effect is even stronger when increasing the Dark Disk contribution, *i.e.* for large values of the relative velocity ρ_{DD}/ρ_H. Interestingly, a co-rotating Dark Disk can boost the mean significance of a Directional Dark Matter detection. Indeed, for a velocity dispersion $\sigma_{DD} = 85$ km/s and a WIMP mass of 50 GeV/c^2, one can see that for rotation velocity $V_{DD} \leq 140$ km/s and for any relative density, the mean significance obtained is greater than the one obtained in the no Dark Disk case. This enhancement of the significance at low rotation velocities can be explained by the fact that the Dark Disk is a colder structure than the host halo, *i.e.* has a smaller velocity dispersion, implying an even more anisotropic recoil angular distribution. Hence, a co-rotating Dark Disk will not necessarily degrade the expected performance of directional detection. Of course, for a perfectly co-rotating Dark Disk ($V_{DD} = v_\odot = 220$ km/s), whatever the velocity dispersion, the recoil angular distribution induced by the Dark Disk is necessarily isotropic. Only the contribution of the Dark Disk to the total number of events will change.

Figure 1 (right) presents the mean significance as a function of V_{DD}. For any value of the velocity dispersion



FIG. 2: Relative asymmetry $\mathscr{A}$ in the plane (V_{DD}, σ_v^{DD}) for a relative density $\rho_{DD}/\rho_H = 1/3$ (left) and $\rho_{DD}/\rho_H = 1$ (right). The solid and dashed lines correspond to isocontours of a relative asymmetry equal to ±1% and ±5% respectively. These studies have been done for 50 GeV/c^2 WIMP mass.

σ_{DD}, the mean significance is continuously decreasing with the rotation velocity of the Dark Disk. However, the case $\sigma_{DD} = 35$ km/s (red solid line) tends to the no Dark Disk limit due to the fact that the contribution to the total number of WIMP events from the Dark Disk falls quickly to zero for $V_{DD} > 140$ km/s. This also explains the rapid decrease of the signficance enhancement in the range $0-100$ km/s. For larger velocity dispersions, the mean significance does not tend to the no Dark Disk limit as the contribution of the Dark Disk to the total number of events remains non negligible. Interestingly, one may note that the range of the values of V_{DD} inducing an enhancement of the significance depends strongly on σ_{DD}. Indeed, for large values of the velocity dispersion, the Dark Matter signal gets closer to an isotropic distribution. This constatation implies that lower is the velocity dispersion, larger is the range in values of V_{DD} allowing for a boost of the directional signature. As a conclusion, larger is the velocity dispersion of the Dark Disk, weaker is the directional discovery significance, except for the case of $\sigma_{DD} = 35$ km/s as discussed above. However, note that the value of $\sigma_{DD} = 141$ km/s is extremely large with respect to the recent results from N-Body simulations. Hence, for a 50 GeV/c^2 WIMP mass, one could expect that a co-rotating Dark Disk could have a positive, though small (∼ 10%), effect on the directional detection of Dark Matter.

For completeness, we studied the evolution of the modifications of the angular distribution $dR/d\Omega_r$ for various Dark Disk parameter values. For this purpose, we defined the relative asymmetry $\mathscr{A}$ as:

$$\mathscr{A} = \frac{\langle\cos\theta\rangle - \langle\cos\theta\rangle_H}{\langle\cos\theta\rangle_H} \qquad (7)$$

where $\langle\cos\theta\rangle_H$ corresponds to the mean recoil deviation obtained in the no Dark Disk case ($\rho_{DD} = 0$). Note that considering the standard halo model and a WIMP of 50 GeV/c^2, we found $\langle\cos\theta\rangle_H \simeq 0.51$. Figure 2 presents the relative asymmetry $\mathscr{A}$ in the plane (V_{DD}, σ_v^{DD}) for a relative density $\rho_{DD}/\rho_H = 1/3$ (left) and $\rho_{DD}/\rho_H = 1$ (right). One may notice that there are three different regions : no effect (the 1% region), a directional discovery enhancement region (low V_{DD}) and a region for which the Dark Disk weakens the directional signature (high V_{DD} together with a high σ_{DD} value). The relative density only affects the amplitude of $\mathscr{A}$, note that the latter spans the range [-18,10] for $\rho_{DD}/\rho_H = 1/3$ and [-40,23] for $\rho_{DD}/\rho_H = 1$. This also affects the area of the no effect region which decreases with increasing value of ρ_{DD}/ρ_H. Interestingly, one can notice that most of the Dark Disk models suggested by N-Body simulations lie in the no effect region. Only extreme, yet unrealistic, Dark Disk models may affect significantly the directional signature.

We evaluate the effect of the Dark Disk contribution to the Dark Matter reach of upcoming directional detectors. Following [26], we compute the directional reach in the (m_χ, σ_p) plane, *i.e* the lower bound of the 3σ discovery region at 90% CL for the two approaches: profile likelihood (red lines) and mean recoil deviation (blue lines). Figure 3 presents the discovery limit in the $(m_\chi, \log_{10}(\sigma_p))$ plane corresponding to two Dark Matter models: standard halo model only (solid lines) and and with an extreme Dark Disk model contribution $\{\rho_{DD}/\rho_h = 1, V_{DD} = 220$ km/s, $\sigma_{DD} = 106$ km/s$\}$ (dashed lines).
The conclusion of this study is twofold. First, we found that for both statistical tests, the effect of an extreme Dark Disk is only mild. Indeed, the directional reach is only degraded by a factor of 3 at high WIMP masses and not affected for light WIMP. Second, we found that the two statistical tests give similar results with a maximal deviation of a few percent. However, it is worth emphasizing that there intepretation differ as the profile



FIG. 3: Discovery limit in the $(m_\chi, \log_{10}(\sigma_p))$ plane corresponding to two Dark Matter models: standard halo model only (solid lines) and and with an extreme Dark Disk model contribution $\{\rho_{DD}/\rho_h = 1$, $V_{DD} = 220$ km/s, $\sigma_{DD} = 106$ km/s$\}$ (dashed lines). We compute the directional reach, *i.e* the lower bound of the 3σ discovery region at 90% CL, for the two approaches: profile likelihood (red lines) and mean recoil deviation (blue lines).

likelihood method favors the background plus signal hypothesis (H_1) whereas the mean recoil deviation method rejects the isotropy hypothesis.

V. CONCLUSION

A co-rotating Dark Disk, as predicted by recent N-Body simulations, might contribute (10%-50%) to the local Dark Matter density, with a potentially dramatic effect on directional detection. In this letter, we have evaluated the effect of Dark Disk model on the discovery potential of upcoming directional detectors. We conclude that, if a co-rotating Dark Disk is present in our Galaxy and has the properties predicted by N-Body simulations [6], the discovery potential of directional detection would be strictly unchanged. Only an extreme, yet unrealistic, Dark Disk model might affect significantly the Dark Matter reach of upcoming directional detectors, by increasing the discovery limit by a factor of three at high WIMP mass ($m_\chi \sim 1000$ GeV/c^2).
Note that even if the impact of Dark Disk contribution to the local Dark Matter distribution only mildly affects the discovery potential of directional detection, it may significantly affect the mass and cross section determination [27]. Indeed, as explained in [12], WIMP events arising from the Dark Disk contribution will induce an excess at low recoil energies which can lower the estimation of the WIMP mass when considering a standard halo model. This highlights the fact that even if a co-rotating Dark Disk is not a threat to the discovery potential of directional detection, it has to be characterized in order to consistently constrain the Dark Matter properties.

[1] R. Teyssier, Astron. Astrophys. **385** (2002) 337
[2] M. Lisanti and D. N. Spergel, arXiv:1105.4166
[3] M. Vogelsberger *et al.*, Mon. Not. Roy. Astron. Soc. **395** (2009) 797
[4] M. Kuhlen, M. Lisanti and D. N. Spergel, arXiv:1202.0007 [astro-ph.GA].
[5] J. Diemand *et al.*, Nature **454** (2008) 735
[6] F. S. Ling, E. Nezri, E. Athanassoula and R. Teyssier, JCAP **1002** (2010) 012
[7] J. I. Read, G. Lake, O. Agertz and V. P. Debattista, MNRAS **389** (2008) 1041-1057
[8] J. I. Read, L. Mayer, A. M. Brooks, F. Governato and G. Lake, MNRAS **397** (2009) 44
[9] C. W. Purcell, J. S. Bullock and M. Kaplinghat, Astrophys. J. **703** (2009) 2275
[10] T. Bruch, A. H. G. Peter, J. Read, L. Baudis and G. Lake, Phys. Lett. B **674** (2009) 250
[11] F. S. Ling, Phys. Rev. D **82** (2010) 023534
[12] A. M. Green, JCAP **1010** (2010) 034
[13] T. Bruch, J. Read, L. Baudis and G. Lake, Astrophys. J. **696** (2009) 920
[14] D. N. Spergel, Phys. Rev. D **37** (1988) 1353
[15] J. Billard, F. Mayet and D. Santos, Phys. Rev. D **82** (2010) 055011
[16] S. Henderson, J. Monroe and P. Fisher, Phys. Rev. D **78** (2008) 015020
[17] C. J. Copi and L. M. Krauss,, Phys. Rev. D **63** (2001) 043507
[18] A. M. Green and B. Morgan, Phys. Rev. D **77** (2008) 027303
[19] C. J. Copi and L. M. Krauss, Phys. Lett. B **461** (1999) 43
[20] B. Morgan, A. M. Green and N. J. C. Spooner, Phys. Rev. D **71** (2005) 103507
[21] B. Morgan and A. M. Green, Phys. Rev. D **72** (2005) 123501
[22] A. M. Green and B. Morgan, Astroparticle Physics 27 (2007) 142
[23] C. J. Copi, L. M. Krauss, D. Simmons-Duffin and S. R. Stroiney, Phys. Rev. D **75** (2007) 023514,
[24] A. M. Green and B. Morgan, Phys. Rev. D **81** (2010) 061301
[25] J. Billard, F. Mayet, J. F. Macias-Perez and D. Santos, Phys. Lett. B **691** (2010) 156-162
[26] J. Billard, F. Mayet and D. Santos, Phys. Rev. D **85** (2012) 035006
[27] J. Billard, F. Mayet and D. Santos, Phys. Rev. D **83** (2011) 075002
[28] S. K. Lee and A. H. G. Peter, JCAP **1204** (2012) 029
[29] D. S. M. Alves, S. E. Hedri and J. G. Wacker, arXiv:1204.5487 [astro-ph.GA].
[30] D. Albornoz Vásquez, G. Bélanger, J. Billard and F. Mayet, Phys. Rev. D **85** (2012) 055023
[31] N. Bozorgnia, G. B. Gelmini and P. Gondolo, arXiv:1111.6361 [astro-ph.CO].
[32] M. S. Alenazi and P. Gondolo, Phys. Rev. D **77** (2008)

043532

[33] R. J. Creswick, S. Nussinov and F. T. Avignone, Astropart. Phys. **35** (2011) 62

[34] N. Bozorgnia, G. B. Gelmini and P. Gondolo, arXiv:1205.2333 [astro-ph.CO].

[35] P. Gondolo, Phys. Rev. D **66** (2002) 103513

[36] M. Lisanti and J. G. Wacker, Phys. Rev. D **81** (2010) 096005

[37] C. M. Bidin, G. Carraro, R. A. Mendez and W. F. van Altena, Astrophys. J. **724** (2010) L122

[38] S. Ahlen *et al.*, Int. J. Mod. Phys. A **25** (2010) 1

[39] K. Miuchi *et al.*, Phys. Lett. B **686** (2010) 11

[40] S. Ahlen *et al.*, Phys. Lett. **B695** (2011) 124-129

[41] D. Santos *et al.*, EAS Publications Series **53** (2012) 25-31

[42] E. Daw *et al.*, Astropart.Phys. **35** (2012) 397-401

[43] J. D. Lewin and P. F. Smith, Astropart. Phys. **6** (1996) 87.

[44] D. Mei and A. Hime, Phys. Rev. D **73** (2006) 053004

[45] G. Cowan, K. Cranmer, E. Gross, O. Vitells, Eur. Phys. J. **C71** (2011) 1554